



Note

Chromatic difference sequences and homomorphisms

Huishan Zhou

Georgia State University, Atlanta, GA 30303-3083, USA

Received 6 December 1988

Revised 22 February 1991

Abstract

Zhou, H., Chromatic difference sequences and homomorphisms, Discrete Mathematics 113 (1993) 285–292.

This note characterizes graphs with the second term of their chromatic difference sequences equal to 1, and gives a class of graphs, called W_n , that are determined by their chromatic difference sequences. It also gives a large class of n -chromatic graphs for which W_n is a homomorphic image. It is proved that the normalized chromatic difference sequence of the categorical product $G \times H$ dominates the normalized chromatic difference sequence of G and of H .

1. Introduction

We only consider finite undirected graphs without loops and multi-edges in this paper. For any graph G , let $V(G)$ and $E(G)$ denote the vertex set and the edge set of G respectively. The *chromatic difference sequence* of a graph G with chromatic number n is defined by $\text{cds}(G) = (a(1), a(2), \dots, a(n))$ where $\sum_{i=1}^t a(i)$ is the number of vertices in a maximum t -colorable induced subgraph of G [1, 2]. The *normalized chromatic difference sequence* of a graph G is defined by $\text{ncds}(G) = \text{cds}(G)/|V(G)|$. A *homomorphism* f from a graph G to a graph H is a mapping $f: V(G) \rightarrow V(H)$ such that $f(x)f(y) \in E(H)$ whenever $xy \in E(G)$. If in addition f is onto $V(H)$ and each edge in H is the image of some edge in G , then f is said to be *onto* and H is called a *homomorphic image* of G . A sequence (x_1, \dots, x_n) is said to *dominate* the sequence (y_1, \dots, y_n) , denoted by

Correspondence to: Huishan Zhou, Department of Mathematics and Computer Science, Georgia State University, University Plaza, Atlanta, GA 30303-3083, USA.

$(x_1, \dots, x_n) \geq^* (y_1, \dots, y_n)$, if $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$, and $\sum_{i=1}^p x_i \geq \sum_{i=1}^p y_i$ for $p = 1, 2, \dots, n-1$. Although domination is usually defined for sequences with the same number of terms. It is natural to extend the definition of domination to two sequences with a different number of nonzero terms. A sequence $x = (x_1, \dots, x_n)$ is said to dominate the sequence $y = (y_1, y_2, \dots, y_m)$ if $n \leq m$, and $x' = (x_1, \dots, x_n, 0, \dots, 0)$, where there are $m-n$ zeros, dominates y . It is also easy to see that when $x_n \neq 0$, $n > m$ then x can not dominate y .

We already know that any sequence with n nonincreasing terms is the cds of a complete n -partite graph [1]. We now consider those sequences with one term equal to 1, and determine which graphs have these sequences as their cds. Theorem 6 gives us an interesting example: the chromatic difference sequence uniquely determines a graph. Finally Theorem 9 gives us a typical example of characterizing the homomorphic image of G from the properties of cds of G . In general a homomorphism from a graph G to (or onto) another graph W need not induce a domination between the ncds of G and W . However under certain conditions a homomorphism from a graph G to another graph W will imply that $\text{ncds}(G)$ dominates $\text{ncds}(H)$. For example, if there exists a homomorphism from G to W and W is vertex-transitive, then $\text{ncds}(G)$ dominates $\text{ncds}(W)$ [2]. Another example involves the Categorical product of two graphs which is defined as follows. The Categorical product $G \times W$ [4] has as vertex set $V(G) \times V(W)$ and has as edge set the edges $(g, w)(g', w')$ where $gg' \in E(G)$ and $ww' \in E(W)$. The projection mapping from $G \times W$ to G is a homomorphism and $\text{ncds}(G \times W)$ dominates $\text{ncds}(G)$ (Theorem 10), from which it is easy to show that $\lim_{k \rightarrow \infty} \text{ncds}(G^k)$ exists, where G^k is recursively defined by $G^1 = G$, $G^k = G^{k-1} \times G$ (Corollary 11). For the Cartesian powers of a graph, we have similar results [5]. The limit behavior of some other graph theoretical parameters of three different graph products were considered in [3].

2. Main results

Lemma 1. Assume that a graph G of chromatic number n has $\text{cds}(G) = (a(1), \dots, a(n))$ with $a(t) = 1$ where $t \leq n$. Then:

- (1) $V(G) = A \cup B$, where A induces a maximum $(t-1)$ -colorable subgraph with $a(1) + \dots + a(t-1)$ vertices and B induces a complete subgraph with $a(t) + \dots + a(n) \leq n$ vertices.
- (2) If A_1, A_2, \dots, A_{t-1} is a $t-1$ coloring of A , then for any $y \in B$, there exists at least one edge between y and each A_i ($i = 1, 2, \dots, t-1$).
- (3) $a(s) \leq t$ for $s = t+1, t+2, \dots, n$.

Remark. If $t = 1$, then $A = \emptyset$. Hence G is a complete graph with n vertices.

Proof. (1) Let A be the vertex set of a maximum $(t-1)$ -colorable subgraph. Then $|A| = a(1) + \dots + a(t-1)$. Let A_1, \dots, A_{t-1} be a $t-1$ coloring of A and

let $B = V(G) - A$. Thus B induces a complete subgraph with $a(t) + \cdots + a(n) \leq n$ vertices, for otherwise $a(t) \geq 2$.

(2) Now if for $y_0 \in B$ and some $i \in \{1, \dots, t-1\}$, there is no edge between y_0 and A_i , then $A_1 \cup \cdots \cup A_{i-1} \cup [A_i \cup \{y_0\}] \cup \cdots \cup A_{t-1}$ induces a $(t-1)$ -colorable subgraph of G with $a(1) + \cdots + a(t-1) + 1$ vertices, a contradiction.

(3) Suppose, otherwise, that $a(j) > t$ for some $j \in \{t+1, t+2, \dots, n\}$. Since $a(i) > 0$ for any $i \in \{1, 2, \dots, n\}$, we have $\sum_{i=t}^n a(i) = \sum_{i=t, i+j}^n a(i) + a(j) > n - t + t = n$, a contradiction. \square

Lemma 2. Assume that a graph G of chromatic number n has $\text{cds}(G) = (a(1), \dots, a(n))$ with $a(2) = 1$. Then G and its cds is either in Case 1 or in Case 2.

Case 1: (1) $a(3) = \cdots = a(n) = 1$; and

(2) $V(G) = A \cup B$ where $|A| = a(1) - 1$, $|B| = n$, A is an independent set and B induces a complete subgraph. There exists a vertex x_0 in B such that there is no edge between this vertex and A but for all other vertices x of B there exists at least one edge between x and A .

Case 2: (1) every one of $a(3), \dots, a(n)$ is equal to 1, except one, say $a(t)$ ($t \in \{3, 4, \dots, n\}$) which is equal to 2;

(2) $V(G) = A \cup B$, where A is an independent set with $a(1)$ vertices, B induces a complete subgraph with n vertices, for any $y \in B$ there is at least one edge between y and A ; and

(3) if $a(t) = 2$ ($t \in \{3, 4, \dots, n\}$), then there exists a subset $B_1 \subseteq B$, $|B_1| = t$ such that:

(3.1) for any $x \in A$, there exists a $y \in B_1$ such that there is no edge between x and y ; and

(3.2) for any $t \in B_1$, there exists an $x \in A$ such that there is no edge between x and y .

Proof. By Lemma 1, $V(G) = A^* \cup B^*$ where A^* is a maximum independent set with $a(1)$ vertices, B^* induces a complete subgraph with $a(2) + \cdots + a(n)$ vertices, and for any $y \in B^*$ there exists at least one edge between y and A^* .

Since $a(2) + \cdots + a(n) \leq n$, and $a(i) \geq 1$, for $i = 1, \dots, n$, then we have either $a(2) + \cdots + a(n) = n - 1$ or $a(2) + \cdots + a(n) = n$.

Case 1: (1) If $a(2) + \cdots + a(n) = n - 1$, then $a(2) = \cdots = a(n) = 1$ and $|B^*| = n - 1$.

(2) We color B^* by $n - 1$ different colors. If for any $x \in A^*$, there exists a $y \in B^*$ such that there is no edge between x and y , then we can color x by the color of y . Thus G is $(n - 1)$ -colorable, a contradiction. Therefore there exists an $x_0 \in A^*$ which is joined to every y in B^* . Let $A = A^* - \{x_0\}$, $B = B^* \cup \{x_0\}$.

Case 2: (1) If $a(2) + \cdots + a(n) = n$, then every one of $a(3), \dots, a(n)$ is equal to 1, except one, say $a(t)$ ($t \in \{3, 4, \dots, n\}$) which is equal to 2. ($a(2) = 1$ is given by the assumption.)

(2) Let $A = A^*$, $B = B^*$.

(3) Assume that $a(t) = 2$ where $t \in \{3, 4, \dots, n\}$. By the definition of cds, a maximum $(t-1)$ -colorable subgraph has $a(1) + t - 2$ vertices. The maximum t -colorable subgraph has $a(1) + t$ vertices. Let H be the maximum t -colorable subgraph. Let $B_1 = B \cap V(H)$. If $|B_1| > t$, then H cannot be t -colorable. If $|B_1| < t$, then $|V(H)| < a(1) + t$, hence H is not the maximum t -colorable subgraph. Therefore $|B_1| = t$, and $V(H) = A \cup B_1$.

(3.1) If for some $x_0 \in A$ we have that $x_0 y \in E(G)$ for every $y \in B_1$, then the subgraph of H induced by $\{x_0\} \cup B_1$ is a $(t+1)$ -complete subgraph. This contradicts the fact that $\chi(H) = t$.

(3.2) Now suppose that $y_0 \in B_1$ with $y_0 x \in E(G)$ for every $x \in A$. Consider the subgraph $H - y_0$. The $t-1$ vertices of $B_1 - y_0$ can be colored by $t-1$ different colors. For any $x \in A$, by (3.1) there exists a $y \in B_1$ such that $xy \notin E(G)$. y cannot be y_0 , therefore we can color x by the color of y . Thus $H - y_0$, which has $a(1) + t - 1$ vertices, is $(t-1)$ -colorable. This contradicts the fact that $a(1) + \dots + a(t-1) = a(1) + t - 2$. \square

Corollary 3. (1) Any sequence $r = (r_1, r_2, \dots, r_n)$ with $r_2 = 1$, some $r_t > 2$ for $t \in \{3, \dots, n\}$ can not be the chromatic difference sequence of some graph.

(2) Any sequence $r = (r_1, r_2, \dots, r_n)$ with $r_2 = 1$, $r_{t_1} = r_{t_2} = 2$ for some $t_1, t_2 \in \{3, \dots, n\}$ can not be the chromatic difference sequence of some graph.

This corollary can also be derived from [1] as mentioned by the referee.

Definition 4. Let W_n be a graph with $2n$ vertices: $a'_1, a'_2, \dots, a'_n, a_1, a_2, \dots, a_n$. The vertices a'_1, a'_2, \dots, a'_n induce a complete graph K_n and the set of vertices $\{a_1, a_2, \dots, a_n\}$ form an independent set. There are edges between a'_i and a_j ($i = 1, 2, \dots, n; j = 1, 2, \dots, n; i \neq j$), no edge between a_i and a'_i ($i = 1, \dots, n$).

Lemma 5 is not hard to prove.

Lemma 5. $\text{cds}(W_n) = (n, 1, 1, \dots, 1, 2)$ with n terms.

Theorem 6. A graph G is isomorphic to W_n if and only if $\text{cds}(G) = (n, 1, \dots, 1, 2)$ with n terms.

Proof. Obviously two isomorphic graphs have the same cds. Thus $\text{cds}(G) = \text{cds}(W_n) = (n, 1, \dots, 1, 2)$ with n terms by Lemma 5 if G is isomorphic to W_n . Conversely if $\text{cds}(G) = (n, 1, \dots, 1, 2)$ with n terms, then we shall prove that G is isomorphic to W_n . By Case 2 of Lemma 2, $t = n$, $V(G) = A \cup B$, where A is an independent set with n vertices, B induces a complete subgraph with n vertices. For any $x \in A$, there exists $y \in B$ such that there is no edge between x and y . For any $y \in B$, there exists $x \in A$ such that there is no edge between y and x .

Claim 1. *For any $y \in B$, there exists exactly one $x \in A$ such that there is no edge between y and x .*

Proof. Suppose, otherwise, that for $y_0 \in B$, there are two $x_1, x_2 \in A$ such that there is no edge from y_0 to x_1 and x_2 . For any $x \in A - \{x_1, x_2\}$ take $\tau(x) \in B$ such that there is no edge between x and $\tau(x)$. Let $B_1 = \{\tau(x) : x \in A - \{x_1, x_2\}\}$. Then $|B_1| \leq n - 2$. Let H be the subgraph induced by $A \cup \{y_0\} \cup B_1$. If $y_0 \notin B_1$, then H can be colored by $|B_1| + 1 \leq n - 1$ colors. But H has $n + |B_1| + 1$ vertices. This contradicts the fact that t -colorable ($t < n$) subgraphs of G can have at most $n + t - 1$ vertices. If $y_0 \in B_1$, then H can be colored by $|B_1| \leq n - 2$ colors. But H has $n + |B_1|$ vertices, which is a contradiction. \square

Claim 2. *For any $x \in A$, there exists exactly one $y \in B$ such that there is no edge between y and x .*

Since for any $y \in B$, there exists exactly one $x \in A$ such that there is no edge between y and x , we must have $n(n - 1)$ edges between B and A . We already know that for any $x \in A$, there exists a $y \in B$ such that there is no edge between x and y , i.e., for any $x \in A$ there is at most $n - 1$ edges between x and B . Now in order that there are $n(n - 1)$ edges between B and A , for any $x \in A$ we must have exactly $n - 1$ edges between x and B . Therefore for any $x \in A$, there is exactly one $y \in B$ such that there is no edge between x and y .

Therefore there is a one to one correspondence between the vertices of A and B . Let $A = \{a_1, a_2, \dots, a_n\}$, $B = \{b_1, \dots, b_n\}$ be such that there is no edge between a_i and b_i ($i = 1, 2, \dots, n$), then there exists an edge between a_i and b_j for any $i = 1, 2, \dots, n$, any $j = 1, 2, \dots, n$, with $i \neq j$. Thus G is isomorphic to W_n . \square (Theorem 6)

Corollary 7. *If $\text{cds}(G) = (t, 1, \dots, 1, 2, 1, \dots, 1)$ where 2 is in the t -th position, then G contains an induced subgraph which is isomorphic to W_t .*

Lemma 8. *Let G be a graph and $\text{cds}(G) = (a(1), a(2), \dots, a(n))$. Then $a(1) \geq a(i)$ for $i = 2, 3, \dots, n$.*

Proof. Suppose $a(1) < a(t)$ for some $t \in \{2, 3, \dots, n\}$. By definition the maximum $(t - 1)$ -colorable induced subgraph of G has $a(1) + \dots + a(t - 1)$ vertices. The maximum t -colorable induced subgraph, say G_t , of G has $a(1) + a(2) + \dots + a(t - 1) + a(t)$ vertices. Let H_1, H_2, \dots, H_t be the color classes of a t -coloring of G_t . Then H_1 is an independent set of G_t , also an independent set of G , hence $|H_1| \leq a(1)$. Thus $|G_t - H_1| \geq a(2) + \dots + a(t - 1) + a(t) > a(2) + \dots + a(t - 1) + a(1)$. $G_t - H_1$ is a $(t - 1)$ -colorable subgraph of G , which contradicts to the fact that the maximum $(t - 1)$ -colorable subgraph has $a(1) + \dots + a(t - 1)$ vertices. \square

Theorem 9. Let G be a graph with $\text{cds}(G) = (a(1), a(2), \dots, a(n))$ and $a(n) > a(i)$ for $i = 2, 3, \dots, n-1$. Then W_n is a homomorphic image of G .

Proof. Let I be a maximum independence set of G and let $H = G - I$. Let A_1, A_2, \dots, A_n be the color classes in an n -coloring of G . Let $I_i = I \cap A_i$ and let $H_i = H \cap A_i$ ($i = 1, \dots, n$) and mapping f by

$$f(x) = a_i \text{ for } \forall x \in I_i; \quad f(x) = a'_i \text{ for } \forall x \in H_i.$$

We shall show that f is a homomorphism from G onto W_n .

Claim 1. f is a homomorphism from H onto the n -complete subgraph induced by $\{a'_1, \dots, a'_n\}$.

Proof. To show that H is not $(n-1)$ -colorable assume the contrary. Let U_i ($i = 1, \dots, n-1$) be the $n-1$ color classes of a $(n-1)$ -coloring of H and let $|U_i| \geq |U_{i+1}|$ ($i = 1, \dots, n-2$). Then I and U_1, \dots, U_{n-2} determine a $(n-1)$ -colorable subgraph of G . Hence

$$a(1) + \dots + a(n-1) \geq a(1) + |U_1| + \dots + |U_{n-2}|,$$

i.e., $a(n) \leq |U_{n-1}|$ and $a(2) + \dots + a(n-1) \geq |U_1| + \dots + |U_{n-2}|$, so

$$(n-2)a(n) \leq (n-2)|U_{n-1}| \leq |U_1| + \dots + |U_{n-2}| \leq a(2) + \dots + a(n-1),$$

i.e. $a(n) \leq a(i)$ holds for at least one i ($i = 2, \dots, n-1$), which is a contradiction. \square

Thus H is n -chromatic since H is n -colorable. Therefore each A_i ($i = 1, 2, \dots, n$) is non-empty. Furthermore any edge in the complete subgraph induced by $\{a'_1, \dots, a'_n\}$ must be the image of edges of H , otherwise H would be $(n-1)$ -colorable.

Claim 2. Each I_i ($i = 1, 2, \dots, n$) is non-empty.

Proof. Suppose without loss of generality that $I_n = \emptyset$. We know that $a(1) = \sum_{i=1}^{n-1} |I_i|$ since I is a maximum independence set. Since $\sum_{i=1}^{n-1} a(i) \geq \sum_{i=1}^{n-1} |I_i| + \sum_{i=1}^{n-1} |H_i|$, we have $a(2) + \dots + a(n-1) \geq |H_1| + \dots + |H_{n-1}|$. On the other hand, $a(1) + a(2) \geq \sum_{i=1}^{n-1} |I_i| + |H_n|$, hence we have $a(2) \geq |H_n|$. Thus

$$\begin{aligned} \sum_{i=1}^n |H_i| &= a(2) + \dots + a(n-1) + a(n) > a(2) + \dots + a(n-1) + a(2) \\ &\geq \sum_{i=1}^n |I_i|. \end{aligned}$$

a contradiction. Hence $I_n \neq \emptyset$. Similarly $I_i \neq \emptyset$ for $i = 1, 2, \dots, n-1$. \square

Claim 3. Each I_i contains a vertex which is adjacent to each H_j ($j \in \{1, 2, \dots, n\}, j \neq i$).



Fig. 1.

Suppose, without loss of generality, that I_n does not contain a vertex which is adjacent to each H_i ($i = 1, \dots, n-1$). Let I_n denote those vertices in I_n which are adjacent to no vertex in H_i . Clearly $I_n = I_{n_1} \cup I_{n_2} \cup \dots \cup I_{n_{n-1}}$. Each $A_i \cup I_n = I_i \cup H_i \cup I_n$ ($i = 1, 2, \dots, n-1$) is an independence set. Thus $a(1) + \dots + a(n-1) \geq |V(G)| - |H_n|$, consequently $a(n) \leq |H_n|$. Since I and H_n form a 2-colorable subgraph of G , we have $a(2) \geq |H_n|$. Thus $a(2) \geq a(n)$, a contradiction. \square (Theorem 9)

The converse of this theorem is not true. For example, we may take G as in Fig. 1.

Theorem 10. *Let G and H be graphs. Then*

$$\text{ncds}(G \times H) \geq^* \text{ncds}(G) \quad \text{and} \quad \text{ncds}(G \times H) \geq^* \text{ncds}(H).$$

Proof. We only need to prove that $\text{ncds}(G \times H) \geq^* \text{ncds}(G)$. Let $\text{cds}(G) = (a(1), a(2), \dots, a(t), \dots, a(n))$, $\alpha(t) = a(1) + \dots + a(t)$ ($t = 1, 2, \dots, n$); $\text{cds}(G \times H) = (b(1), b(2), \dots, b(t), \dots, b(n))$, $\beta(t) = b(1) + \dots + b(t)$. Let $I_t = \{x_1, \dots, x_{\alpha(t)}\}$ ($1 \leq t \leq n$) be the vertices of the maximum t -colorable subgraph of G . Then $\{x_1\} \times V(H) \cup \{x_2\} \times V(H) \cup \dots \cup \{x_{\alpha(t)}\} \times V(H)$ are the vertices of a t -colorable subgraph of $G \times H$. Thus

$$\beta(t) \geq \sum_{i=1}^{\alpha(t)} |\{x_i\} \times V(H)| = \alpha(t) |V(H)| \quad (t = 1, \dots, n-1),$$

and

$$\beta(n) = \sum_{i=1}^{\alpha(n)} |\{x_i\} \times V(H)| = |V(G)| |V(H)|.$$

Therefore

$$\frac{\beta(t)}{|V(G \times H)|} \geq \frac{\alpha(t) |V(H)|}{|V(G)| |V(H)|} = \frac{\alpha(t)}{|V(G)|} \quad (t = 1, 2, \dots, n-1),$$

and

$$\frac{\beta(n)}{|V(G \times H)|} = 1 = \frac{\alpha(n)}{|V(G)|}.$$

i.e., $\text{ncds}(G \times H) \geq^* \text{ncds}(G)$. \square

Corollary 11. *Let G be a graph, and G^k be recursively defined by $G^1 = G$, $G^k = G^{k-1} \times G$. Then $\lim_{k \rightarrow \infty} \text{ncds}(G^k)$ exists.*

Acknowledgments

The author wishes to thank referees for corrections and suggestions.

References

- [1] M.O. Albertson and D.M. Berman, The chromatic difference sequence of a graph, *J. Combin. Theory Ser. B* 29 (1980) 1–12.
- [2] M.O. Albertson and K.L. Collins, Homomorphism of 3-chromatic graphs, *Discrete Math.* 54 (1985) 127–132.
- [3] P. Hell and F.S. Roberts, Analogues of the Shannon capacity of a graph, *Ann. Discrete Math.* 12 (1982) 155–168.
- [4] D.J. Miller, The Categorical product of graphs, *Canad. J. Math.* 20 (1968) 1511–1521.
- [5] H. Zhou, The chromatic difference sequence of Cartesian product of graphs, *Discrete Math.* 90 (1991) 297–311.